

**UDC 55**

**CODEN – GEOME 2**

**ISSN 0352 – 1206**

# **GEOLOGICA MACEDONICA**

<i>Geologica Macedonica</i>	Год.	24	Број	1	стр.	1–74	Штип	2010
<i>Geologica Macedonica</i>	Vol.	No		pp.			Štip	

<i>Geologica Macedonia</i>	Год.	24	Број	1	стр.	1–74	Штип	2010
<i>Geologica Macedonia</i>	Vol.	No		pp.			Štip	

## GEOLOGICA MACEDONICA

Published by: – Издава:

The "Goce Delčev" University, Faculty of Natural and Technical Sciences, Štip, Republic of Macedonia  
Универзитет „Гоце Делчев“, Факултет за природни и технички науки, Штип, Република Македонија

### EDITORIAL BOARD

**Todor Serafimovski** (R. Macedonia, *Editor in Chief*), **Prof. Blažo** (R. Macedonia, *Editor*), David Alderton (UK), Tadej Dolenc (R. Slovenia), Ivan Zagorchev (R. Bulgaria), Wolfgang Todt (Germany), acad. Nikolay S. Bortnikov (Russia), Clark Burchfiel (USA), Thierry Augé (France), Todor Delipetrov (R. Macedonia), Vlado Bermanec (Croatia), Milorad Jovanovski (R. Macedonia), Spomenko Mihajlović (Serbia), Dragan Milovanović (Serbia), Dejan Prelević (Germany), Albrecht von Quadt (Switzerland)

### УРЕДУВАЧКИ ОДБОР

**Тодор Серафимовски** (Р. Македонија, *главен уредник*), **Блажо Боев** (Р. Македонија, *уредник*), Дејвид Олдертон (В. Британија), Тадеј Доленец (Р. Словенија), Иван Загорчев (Р. Бугарија), Волфганг Тод (Германија), акад. Николай С. Бортников (Русија), Кларк Барв菲尔 (САД), Тиери Оже (Франција), Тодор Делипетров (Р. Македонија), Владо Берманец (Хрватска), Милорад Јовановски (Р. Македонија), Споменко Михајловић (Србија), Драган Миловановић (Србија), Дејан Прелевић (Германија), Албрехт вон Квад (Швајцарија)

Language editor	Лектура
<b>Marijana Kroteva</b>	<b>Маријана Кротева</b>
(English)	(англиски)

<b>Georgi Georgievski, Ph. D.</b>	<b>д-р Георги Георгиевски</b>
(Macedonian)	(македонски)

Technical editor	Технички уредник
<b>Blagoja Bogatinoski</b>	<b>Благоја Богатиноски</b>
Proof-reader	Коректор
<b>Alena Georgievska</b>	<b>Алена Георгиевска</b>

Address	Адреса
<b>GEOLOGICA MACEDONICA</b>	<b>GEOLOGICA MACEDONICA</b>
<b>EDITORIAL BOARD</b>	<b>РЕДАКЦИЈА</b>
Faculty of Natural and Technical Sciences	Факултет за природни и технички науки
P. O. Box 96	пошт. фах 96
MK-2000 Štip, Republic of Macedonia	MK-2000 Штип, Република Македонија
Tel. ++ 389 032 550 575	Тел. 032 550 575
E-mail: <a href="mailto:todor.serafimovski@ugd.edu.mk">todor.serafimovski@ugd.edu.mk</a>	

400 copies	Тираж: 400
Published yearly	Излегува еднаш годишно
Printed by:	Печати:
2 <sup>ri</sup> Avgust – Štip	2 <sup>ри</sup> Август – Штип
Price: 500 den	Цена: 500 ден.
The edition was published in December 2010	Бројот е отпечатен во декември 2010

Photo on the cover:

Argillitic alteration, Kadiica, Republic of Macedonia

На корицата:

Аргилитска алтерација, Кадица, Република Македонија

<i>Geologica Macedonica</i>	Год.	24	Број	1	стр.	1–74	Штип	2010
<i>Geologica Macedonica</i>	Vol.	No		pp.			Štip	

## СОДРЖИНА

**Споменко Ј. Михајловиќ, Руди Чоп, Паоло Паланцио**

Структура на големите магнетни бури ..... 1–12

**Марјан Делипетрев, Жан Л. Расон, Благица Донева, Тодор Делипетров**

Мрежа на мерни станици и тектонска реонизација на Република Македонија ..... 13–21

**Милорад Јовановски, Азра Шпаго, Игор Пешевски**

Дијапазон на вредности на инженерско-геолошки карактеристики на некои карбонатни карпести комплекси од балканскиот полуостров ..... 23–30

**Гоше Петров, Виолета Стојанова, Војо, Мирчовски, Андреј Шмуц, Ѓорѓи Димов**

Тектонска еволуција на палеогените басени во Република Македонија ..... 31–37

**Тодор Серафимовски, Горан Тасев, Крсто. Блажев, Александар Волков**

Главните Алписки структури и Си-порфирска минерализација во Српско-Македонскиот масив ..... 39–48

**Тена Шијакова-Иванова, Весна Амбаркова, Vassiliki Topitsoglou, Весна Панева-Зајкова**

Зависност помеѓу концентрацијата на флуор и останатите елементи во некои геотермални води во Република Македонија ..... 49–52

**Снежана Димовска, Трајче Ставилов, Роберт Шајн**

Определување на активноста на  $^{40}\text{K}$  и вкупната бета активност во почвата од Кавадарци и неговата околина ..... 53–62

**Сабина Стрмиќ Палинкаш, Сибила Боројевиќ Шоштариќ, Ладислав Палинкаш,**

**Золтан Печкај, Блажо Боев, Владимир Берманец**

Гасно-течни инклузии и одредување на староста според методот K/Ar на Au-Sb-As-Tl наоѓалиштето Алшар, Македонија ..... 63–71

**Упатство за авторите**

73–74

<i>Geologica Macedonica</i>	Год.	24	Број	1	стр.	1–74	Штип	2010
<i>Geologica Macedonica</i>	Vol.		No		pp.		Štip	

## TABLE OF CONTENTS

<b>Spomenko J. Mihajlović, Rudi Čop, Paolo Palangio</b> The structure of the big magnetic storms.....	1–12
<b>Marjan Delipetrev, Jean L. Rasson, Blagica Doneva, Todor Delipetrov</b> Net of repeat stations and tectonic regionalization of the Republic of Macedonia .....	13–21
<b>Milorad Jovanovski, Azra Špago, Igor Peševski</b> Range of engineering-geological properties for some carbonate rock complexes from Balkan Peninsula.....	23–30
<b>Goše Petrov, Violeta Stojanova, Vojo Mirčovski, Andrej Šmuc, Đordi Dimov</b> Tectonics evolution of the paleogene basins in the Republic of Macedonia.....	31–37
<b>Todor Serafimovski, Goran Tasev, Krsto Blažev, Aleksandr Volkov</b> Major alpine structures and Cu-porphyry mineralization in the Serbo-Macedonian massif.....	39–48
<b>Tena Šijakova-Ivanova, Vesna Ambarkova, Vassiliki Topitsogloy, Vesna Paneva-Zajkova</b> Fluoride content and dependence on other elements in some geothermal waters in Republic of Macedonia .....	49–52
<b>Snežana Dimovska, Trajče Stafilov, Robert Šajn</b> Determination of activity concentration of $^{40}\text{K}$ and gross beta activity in soil from Kavadarci and its environs .....	53–62
<b>Sabina Strmić Palinkaš, Sibila Borojević Šoštarić, Ladislav Palinkaš, Zoltan Pecskay, Blažo Boev, Vladimir Bermanec</b> Fluid inclusions and K/Ar dating of the Allšar Au-Sb-As-Tl mineral deposit, Macedonia .....	63–71
<b>Instructions to authors.....</b>	73–74

## TECTONICS EVOLUTION OF THE PALEOGENE BASINS IN THE REPUBLIC OF MACEDONIA

Goša Petrov<sup>1</sup>, Violeta Stojanova<sup>1</sup>, Vojo Mirčovski<sup>1</sup>, Andrej Šmuc<sup>2</sup>, Đordž Dimov<sup>3</sup>

<sup>1</sup>Faculty of Natural and Technical Sciences, "Goce Delčev" University,  
Goce Delčev 89, MK-2000, Štip, Republic of Macedonia

<sup>2</sup>Faculty of Natural and Technical Sciences, Geology Department, University in Ljubljana  
Privoz 11, 1000 Ljubljana, Slovenia

<sup>3</sup>Faculty of Computer Sciences, "Goce Delčev" University, ,  
Tošo Arsov 14, MK-2000 Štip, Republic of Macedonia  
gose.petrov@ugd.edu.mk // violeta.stojanova@ugd.edu.mk


**A b s t r a c t:** Paleogene sediments on the territory of the Republic of Macedonia are widespread, especially in its central and eastern parts, i.e. in the Vardar zone and Serbo-Macedonian massif (Kraisthida zone). After Laramide orogeny compression, in expansion conditions, along the gravitational faults, continental trenches were created on these spaces, with general direction NW–SE, in which paleogene molassa sediments were accumulated. In the Vardar zone, continental trenches are extended to the north to Skopska Crna Gora mountain (north of Skopje), which includes the Skopje–Kumanovo basin, Ovče Pole basin, Tikveš basin, Gevgelija–Strumica basin and Valandovo basin. In Kraisthida zone continental trenches stretches along the border with Bulgaria and it includes Deve Bair basin and Delčeva basin. Bearing in mind the different thickness of the allocated Paleogene lithozones in the separate basins, and the fact that in some basins lithozones absent, it can be assumed that their spreading along trenches were segmented by cross faults with NE–SW direction. Cross faults are mainly contemporaneous faults.

**Key words:** Paleogene; molassa sediments; continental trench; contemporaneous fault

## INTRODUCTION

After the final closure of the Vardar Ocean and finishing processes of collision with Laramide orogeny phase (Latest Cretaceous–Paleocene), the southern parts of the Balkan Peninsula (the territory of Macedonia, southern Bulgaria, northern Greece and eastern Albania), in the Eocene–Oligocene period, became in the conditions of expansion in the south Balkan extension region (SBER) (Dumurdžanov et al. 2005). This extension system belongs to a much wider area, separated as Aegean extension region (Sengor et al. 2004) (Fig. 1).

Besides the intensive tectonics of Paleogene sediments with Pyrenian orogeny and Savian orogeny compression (folding, faulting and shelling), lithostratigraphy of these sediments is relatively simple. Along the edge parts of the basin, sediments are represented by coarse conglomerates and sandstones, while internal parts are represented by finegranular sediments, marlstone, clay and others. They are transgressive through the older formations, from Precambrian to Mesozoic.



**Fig. 1.** Simplified tectonic map of the Eastern Mediterranean region showing Southern Balkan Extensional Region (SBER) – horizontal lines) in relation to selected tectonic features. In Balkan region, position of the volcanic arcs of Eocene–Oligocene age (Eo-Olig); Miocene (Mio), and Pliocene to Recent (Active); are shown. The location of Macedonia is outlined. KF – Kefalonia fault zone; NAF – North Anadolska fault zone (taken from Dumurdžanov et al. 2005)


In all these basins, Paleogene sediments have been identified as Upper Eocene–Priabonian and Lower Oligocene (Pavlović 1926; Темкова 1958, 1967; Mitrović et al. 1990; Džuranov et al. 1999;

Туњева 2000). The total thickness of Paleogene sediments (Upper Eocene–Lower Oligocene) is evaluated at about 3 000–3 500 m.

### STRATIGRAPHY OF PALEOGEN IN THE R. MACEDONIA

According to the spatial expansion, Paleogene in R. Macedonia can be divided into 4 main (larger) basins and a few isolated blocks, usually located along the shells with NW-SE orientation. These are: Tikveš, Ovče Pole, Skopje–Kumanovo

and Delčeve basin as the main basins and Deve Bair, Dedeli–Rabrovo (near Valandovo) and Illovica–Štuka in Strumica valley, as isolated blocks (Fig. 2).




**Fig. 2.** Distribution of Paleogene sediments in Macedonia  
1 – Tikveš basin, 2 – Ovče Pole basin, 3 – Skopje–Kumanovo basin, 4 – Delčeve basin, 5 – Valandovo–Gevgelija basin,  
6 – Strumica basin, 7 – Deve Bair basin

With lithostratigraphical studies to Paleogene sediments in the basins of R. Macedonia, five superpositional lithostratigraphical units (litozones) were separated: basal lithozone, low flysch lithozone, lithozone of yellow sandstones, upper flysch lithozone and Oligocene lithozone (Fig. 3).

Basal lithozone ( $^1E_3$ ) starts with basal conglomerates and sandstones, clay stone and kalkarenits (all in red and purple color) lay above. This zone is presented in all Paleogene basins on the territory of the R. Macedonia. Thickness of this lithozone varies from 350–700 m. Paleontological is docu-

mented in the Tikveš basin, with large macrofauna, represented by Molusca and Anthozoa (Maksmović et al. 1954).

Lower flysch lithozone ( $^2E_3$ ) is a thick about 1100 m. It is developed in Tikveš, Ovče Pole and Deve – Bair basin with rhythmically replacement of red and gray conglomerates, sandstones, claystones and marlstone and alevrolites. In Deve Bair basin, in the rhythms of the zone volcanogenetic sediments appear. In the upper stratigraphic levels Priabonian snails, shells, coral and macroforaminifer fauna are determined (Темкова 1958).



**Fig. 3.** Correlation of Paleogene basins in the Republic of Macedonia

1 – lower oligocene sediments, 2 – upper flysch lithozone, 3 – lithozone of yellow sandstones, 4 – lower flysch lithozone,  
5 – basal lithozone, 6 – paleorelief

Lithozone of yellow sandstones ( ${}^3E_3$ ) is thick from 600 to 1000 m. It is present in Tikveš and Ovče Pole basin, where vertical and lateral passes in the top flysch lithozone, while in Skopje–Kumanovo basin sediments that belong to this lithozone were moved out into the sediments with Oligocene age. Lithozone of yellow sandstones is determined as Priabonian with large foraminifer fauna, snails, corals, shells and more (Темкова 1967).

Upper flysch lithozone ( ${}^4E_3$ ) is thick from 1000 – 2000 m. The greatest thickness (about 2000 m) is presented in Tikveš basin (based on data from drilling). Characteristic for this lithozone is its gray color. In the upper flysch lithozone gray sandstone, marls and claystones are present and on the top are limestones, with obvious gradation, convolution and lamination.

All levels of the lithozone are rich in snails, shells and foraminifer fauna, while in the upper limestone layers are determined: *Nummulites fabiani*, *Nummulites incrassatus*, *Nummulites budensis*, *Operculina alpina*, *Discocyclina angastae*

(Maksimović et al. 1954, Ракичевиќ et al. 1976) and microforaminifer fauna of Priabonian age (Džuranov et al. 1999).

Oligocene sediments are preserved with thickness from 200–300 m in some parts of Ovče Pole basin (in the vicinity of Kočani) and in Skopje –Kumanovo basin. This lithozone begins with ridge gray limestones, with typical marine fauna the sandstones are above them and on the top ends with clay stones.

Limestones have typical shallow character. In the lower part they are in breccious, and in the upper part move into corale limestone. The age of limestone is problematic, because despite Oligocene types of Nummulti (*Nummulites intermedius* – typical oligocene kind), Anthozoa, Gasteropoda, Bivalvia and Echinodermata (Karajovanović et al. 1975), in fossil association occur species from Eocene to Oligocene. But according to stratigraphy position, lithologic composition and discovered fauna, the age of these sediments is determined as the lower Oligocene.

## TECTONICS EVOLUTION OF PALEOGEN AND DEVELOPMENT OF ACUMULATION BASIN IN THE R. MACEDONIA

In terms of expansion and a general trend of segmentation in the wider region of the Balkan Peninsula, wide areas of the territory of the Republic Macedonia, primarily the area of the Vardar zone and Serbo-Macedonian massif (Kraisthida zone) and border areas with Albania (near Debar), during Upper Eocene were covered with differential vertical movements. Such processes of extension and vertical tectonics conditioned the formation of trenches, along which gradually entering marine waters from the south and southeast to the north and northwest. Today preserved Eocene–Oligocene masses with its geological position, extension and orientation suggest that the formation of trenches was controlled by the gravitational faults with a dominant direction NW–SE. Faults in this direction, often represent reactivated older faults, located in the border, very mobile, parts of the Vardar zone with Pelagonian massif (from west), Serbo-Macedonian massif (eastern side) and the newly formed fault lines in the Vardar zone and Serbo-Macedonian massif of the same orientation (NW–SE).

With the intensive transgression, because of the spreading of the sedimentation basin, sedimentation from the trenches expanded and spread to most of the Vardar zone. Towards north reach the terrains northeast of Skopje, and to the east covered the entire central and eastern Vardar subzone, and most of the Serbo – Macedonian massif (and Kraisthida zone). In that way, very wide sedimentary basin that existed in the Upper Eocene was formed.

The accumulation of Upper Eocene sediments began with continental terrigenous sediments, with red and red-violet color, presented by coarse conglomerates, sandstones and clay stones and thin layers of gray kalkarenites. With the gradual deep-

ening of the area, sedimentation become sea type. Vertical sinking of the terrains was the most expressive in Tikveš basin, where sediments are formed by mass with thickness 3 000–3 500 m and in Ovče Pole basin to 3 000 m (Fig. 3).

Paleogene sedimentary masses from Tikveš, Ovče Pole and Skopje–Kumanovo basin presented, more or less, a whole, in places covered by Neogene or Quarter deposit or divided by a narrow neotectonics horst. The general orientation of this mass is NW–SE, conditioned by postlaramie faults segmentation of the area, with which trenches were formed and after which entering Eocene transgression with accumulation of sediments. Most of these fault structures were renewed in posteocene period, and because of that mass maintaining the same orientation. Small deviations in the extension of mass in the WNW–ESE direction are caused by faults of neotectonic stage.

After the forming of Late Eocene–Oligocene sediment mass, in the Lower Oligocene, it was covered with Pyrenian compression, and at the end of Oligocene and Lower Miocene with Savian compression. During these orogeny phases, intensive folding, faulting and sheathing was made towards the west. As a result, widespread Paleogene mass was segmented in a large number of partially or completely isolated blocks. Compression oriented east–west, conditioned folding of Paleogene sediments in discrete anticline and syncline form, often with an orientation NW–SE to N–S. With this compression, parallel with folding, creation of numerous faults and shells with NW–SE to N–S orientation starts.

## RESULTS AND DISCUSSION

More conclusions can be done if analyzing the horizontal distribution of Upper Eocene–Lower Oligocene sediments (separated in 5 superpositional lithozone) in individual sedimentary basins, and analysis of the thickness of individual lithozone. Namely, it was mentioned that basal lithozone ( ${}^1E_3$ ) is present in all Paleogene basins, suggesting that in this period sedimentation took place

in relatively calm tectonic conditions. The thickness of the basal lithozone is the largest in Tikveš and Gevgelija–Valandovo basin, which means that in such a relatively calm tectonic conditions southern part of Paleogene sedimentary basin in the territory of the R. Macedonia sinking faster.

During the deposition of the lower flysch lithozone ( ${}^2E_3$ ), differentiated vertical movements

in sedimentary basin took place. This lithozone is absent in Skopje–Kumanovo, Delčevo, Gevgelija–Valandovo and Strumica basin, which means that these regions were land in that time. This suggests that in addition to faults with spreading NW–SE (after formation of Paleogene trenches) existed transverse fault lines in NE–SW direction (Fig. 4), after which becomes differentiated vertical rising (or sinking).


In the Vardar zone, lower flysch lithozone has larger thickness in the Ovče Pole basin than in Tikveš basin, and it can be concluded that along the transverse contemporaneous fault, Ovče Pole basin is sinking faster. In border areas of Bulgaria (in Kraisthide zone) in the northern trench (Deve Bair basin) sinking was more intense compared to the southern part (Delčevo basin).

Lithozone of yellow sandstones ( ${}^3E_3$ ), as the lower flysch lithozone is not continuous in all Paleogene basins. It is absent in Delčevo basin, Gevgelija–Valandovo basin and in Strumica basin. The trench in the Vardar zone has the greatest thickness

in the Skopje–Kumanovo basin, and along the transverse fault sank relatively faster compared to Ovče Pole and Tikveš basins. In the Serbo–Macedonian basin lithozone of yellow sandstones is not present, meaning that these areas were land.

In the final phase of Paleogene sedimentation, at the time of deposition of upper flysch lithozone ( ${}^4E_3$ ), the Vardar zone and Serbo–Macedonian massif (Kraisthide zone) were under water. The southern parts of Vardar zone (Tikveš and Ovče Pole basin) sinking more intense compared to the northern parts (Skopje–Kumanovo basin). In Kraisthide zone southern of the trench (Delčevo basin) sinking more intense compared to the north, where the upper flysch lithozone is eroded.

In the Lower Oligocene (Ol) sedimentation took place mainly in the northern parts of continental trench along the Vardar zone. In Skopje–Kumanovo basin sinking was most intense, and in the SE terrains relative rising was present.



**Fig. 4.** Simplified tectonic map of the Paleogene in the Republic of Macedonia  
Paleogene continental trenches: I – in Vardar zone, II – in Serbo–Macedonian massif (Kraisthide zone); 1 – paleogene sediments,  
2 – longitudinal faults, 3 – cross (transverse) faults

## CONCLUSION

On the territory of the Republic of Macedonia Paleogene sedimentary basins are present in two continental trenches formed in extension conditions along the gravitational fault lines with direction NW–SE. Continental trench that was formed in Vardar zone, starts north of Skopje–Kumanovo basin. Towards SE structure expands and includes the Ovče Pole basin, Tikveš basin, Gelgelija–Valandovo basin and Strumica basin. In the border belt with Bulgaria, in the Kraisthida zone, relatively narrower continental trench that covered Deve Bair basin and Delčeve basin was formed.

Besides gravitational edge faults, along which the continental trenches were formed, great signi-

fificance for the course and character of the individual Paleogene sedimentation basins have cross-faults with NE–SW direction. Along these, mainly contemporaneous fault, during the Upper Eocene – Lower Oligocene differentiated vertical movements took place. As a result of such sinking or rising, big differences in thickness of individual lithozone (Fig. 3) appeared. In Tikveš basin and Ovče Pole basin the Paleogene lithostratigraphic columns are complete, while in other Paleogene basins one or two lithozones are missing (lower flysch lithozone and the lithozone of yellow sandstone)

## REFERENCES

- Arsovski M., Dumurdžanov N., 1995: Alpine tectonic evolution of the Vardar zone and its place in the Balkan region. *Geologica Macedonica*, **9**, 1, 15–22,
- Арсовски М., 1997: *Тектоника на Македонија*. Рударско-геолошки факултет, Штип.
- Arsovski M., Dumurdžanov N., Petrov G., 1997: *Manifestations of the Alpine orogen phases in the Vardar zone*. Proceeding, Symposium, Dojran.
- Dumurdžanov N. et al., 2005: *Cenozoik tectonics of Macedonia and its relation to the South Balkan extensional regime*. *Soesphere*, Geological Society of America, p. 1–22.
- Džuranov S., Tuneva V., Dumurdžanov N., 1999: Microforaminifera findings near the village of Čardaklja in the Ovče pole paleogene basin, Republic of Macedonia. *Geologica Macedonica*, **13**, 57–68.
- Јанчевски Ј., 1987: *Класификација на раседниште структури и тенеза, стапости и морфологија со осврт на сеизмичноста на територијата на Македонија*. Докторска дисертација, РГФ, Штип.
- Maksimović B., Sikošek B., Marković O., & Veselinović M., 1954: Geološki sastav i tektonска struktura jednog dela Ovčeg Polja i Tikveša sa paleontološkom dokumentacijom. *Trudovi na Geološki Zavod na NRM*, Sv. 4, Skopje, p. 1–177.
- Карајовановик М. & Хаци-Митрова С., 1975: *Толкувач за Основна Геолошка Карта на СФРЈ*, 1:100 000, лист Титов Велес. Геолошки завод. Скопје.
- Pavlović P., 1926: Priabonski mekuscii sa Ovčeg Polja i Jezevog Brda. *Geol. Anal. Balk. Pol.*, Kn. VIII, 2, Beograd, p. 231–254.
- Петров Г., 2002: *Геодинамика на тектонскиите процеси на Вардарската зона со посебен осврт на Алтиската орогенеза*. Докторска дисертација, РГФ, Штип.
- Петров Г., Стојанова В., Думурџанов Н., Димов Г., 2009: *Корелација на еоценскиите флишини седименти на Македонија и Словенија и влијанието на регионалното палеотектографско проучување на флишините басени*. Факултет за природни и технички науки, Штип.
- Petrović-Mitrović J. et al., 1990: *Paleogene echinoides of Eastern Macedonia*. 13<sup>th</sup> Congress of the Geologists of Jugoslavia, Kn. 1, p. 369–380.
- Ракичевић Т., Думурџанов Н., Петковски П., 1976: *Толкувач за Основна геолошка карта на СФРЈ*, 1:100 000, лист Штип. Геолошки завод, Скопје.
- Sengor et al. 2004: The North Anatolian Fault. A new look. *Ann. Rev. Earth Planet. Sci.* **33**, 1–75.
- Стојанова В., 2008: *Еволуција и стратиграфија на палеотенот на територијата на Република Македонија*. Докторска дисертација, ФРГП, Штип, с. 13–78.
- Темкова В., 1958: Палеонтолошка обработка на фауната во Тиквешкиот басен и околината. *Трудови на Геолошки завод на НРМ*, **6**, с. 93–123, Скопје.
- Темкова В., 1967: Палеонтолошка одредба на макрофауната од палеотените седименти на теренот, лист Штип. Стручен фонд на Геолошки завод на СРМ, Скопје.
- Тунева В., 2000: *Стратиграфија на палеотенот од Овчејолскиот басен според микрофораминиферите*. Магистерска работа. Рударско-геолошки факултет, Штип.

## Резиме

## ТЕКТОНСКА ЕВОЛУЦИЈА НА ПАЛЕОГЕННИТЕ БАСЕНИ ВО РЕПУБЛИКА МАКЕДОНИЈА

Гоше Петров<sup>1</sup>, Виолета Стојанова<sup>1</sup>, Војо Мирчовски<sup>1</sup>, Андреј Шмуц<sup>2</sup>, Ѓорѓи Димов<sup>3</sup>

<sup>1</sup>Универзитет "Гоце Делчев", Факултет за природни и технички науки,  
Гоце Делчев 89, МК – 2000 Штип, Република Македонија

<sup>2</sup>Факултет за природни и технички науки, Оддел за геологија, Универзитет во Љубљана  
Привоз 11, 1000 Љубљана, Словенија

<sup>3</sup>Универзитет "Гоце Делчев", Факултет за информатика,  
Тошо Арсов 14, МК – 2000 Штип, Република Македонија  
gose.petrov@ugd.edu.mk // violeta.stojanova@ugd.edu.mk

**Клучни зборови:** палеоген; моласни седименти; континентален трог; контемпорарен расед.

Палеогените седименти на територијата на Република Македонија се широко распространети, особено во нејзиниот централен и источен дел, односно во Вардарската зона и Српско-Македонскиот масив (Краиштидната зона). По Ларамиската орогена компресија, во услови на екstenзија, на овие простори биле создадени континентални трогови, долж гравитациони раседи со генерален правец на протегање СЗ–ЈИ, во кои се таложеле палеогени моласни седименти.

Во Вардарската зона континенталниот трог на север се простирал до Скопска Црна Гора (северно од Скопје), во кој влегуваат Скопско-Кумановскиот басен, Овчепол-

скиот басен, Тиквешкиот басен, Гевгелиско-Валандовскиот басен и Струмичкиот басен. Во Краиштидната зона континенталниот трог се протега долж границата со Бугарија и во него влегуваат Девебаирскиот басен и Делчевскиот басен.

Имајќи ја предвид различната дебелина на издвоените палеогени литозони во поедините басени, и фактот што во некои басени некои литозони отсуствуваат, може да се претпостави дека троговите долж нивното протегање биле сегментирани со напречни раседи со протегање СИ–ЈЗ. Напречните раседи главно претставуваат контемпорарни раседи.